ACCEPTED
FOR REGISTRATION

5/15/12

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc. ID: 531420

20-0-4
.28% Barricade, PCSCU

GUARANTEED ANALYSIS

Total Nitrogen (N)	20.0%
20.0% Urea nitrogen*	
Soluble Potash (K_2O)	4.0%

Plant foods derived from: Urea, Poly Coated Sulfur Coated Urea and Muriate of Potash.

*5.0% slowly available urea nitrogen from poly coated sulfur coated urea.

390-5040 winpocketsm

WINFIELD™ FERTILIZER WITH Barricade® 0.28%

ACTIVE INGREDIENT:	
Prodiamine [N¹, N²-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine]	0.28%
OTHER INGREDIENTS:	99.72%
TOTAL	100.00%

EPA Reg. No. 9779-363 **1/0126/0** **EPA Est. No. 57131-WI-1**

Controls Crabgrass and Other Weeds Early Spring

For selective preemergence control of grass and broadleaf weeds in:
- **Established turf grasses (excluding golf course putting greens) and lawns**
- **Landscape ornamentals including**
- **Established perennials & wildflower plantings**

Covers Up to 12,500 Square Feet

KEEP OUT OF REACH OF CHILDREN
CAUTION

See the following section, PRECAUTIONARY STATEMENTS, HAZARDS TO HUMANS AND DOMESTIC ANIMALS, for additional information.

Barricade® is a registered trademark of Syngenta.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

If in Eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If Inhaled	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information.

Net Wt. 50 lbs. (22.7 kg)

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE

Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

If material is spilled: Contain and sweep up material of spill for disposal as per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Paper and plastic bags: Nonrefillable container. Do not reuse empty bag. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns, landscape ornamentals, and established perennials & wildflower plantings.

This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in **Maximum Annual Rates** on this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns:

1. Do not graze or feed livestock forage cut from areas treated with this product.
2. Do not use on food producing plants.
3. Do not apply this product through any type of irrigation system.
4. Do not apply aerially.
5. Do not apply to golf course putting greens.
6. Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
7. Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
8. Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.
9. Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
10. Disturbing the herbicide barrier with cultural practices, such as disking may result in reduced weed control.
11. Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bentgrass (Poa annua)are desirable species.
12. Keep this product off sidewalks, patios, driveways, etc. This product contains materials which will stain concrete and painted surfaces. If it comes in contact with a concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate Per Calendar Year By Turf Grass Species[1]		
Turf Species	**Lbs. Product/A**	**Lbs. ai/A**
Bermudagrass[2] Bahiagrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	530	1.5
Buffalograss Kentucky Bluegrass Perennial Ryegrass	353	1.0
Creeping Red Fescue	265	0.75
Creeping Bentgrass	230	0.65

[1] These are the maximum rates per calendar year by species limitations.
[2] May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 183 lbs./A or 4 lbs./1000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

• Do not apply more than 530 lbs. of product per calendar year.
Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the "Maximum Annual Rates" table.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Carpetweed	Itchgrass	Pigweed
Bluegrass, Annual (Poa annua)	Chickweed, Common	Johnsongrass (from seed)	Purslane, Common
Crabgrass (large, smooth)	Chickweed, Mouseear (from seed)	Junglerice	Pusley, Florida
Crowfootgrass	Henbit	Lovegrass	Shepherdspurse(2)
Cupgrass, Woolly	Knotweed	Panicum (Texas, Fall, Browntop)	Speedwell, Persian
Foxtails, Annual	Kochia	Rescuegrass(3)	Spurge, Prostrate
Goosegrass(1)	Lambsquarter, Common	Signalgrass, Broadleaf	Witchgrass
		Sprangletop	Woodsorrel, Yellow (from seed)

(1) In many areas a single application of 230 to 530 lbs./A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 230 to 353 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed those given in the "Maximum Annual Rates" table. Do not exceed the maximum rate for turf grass species listed in the "Maximum Annual Rates" table above.
(2) Applications for this weed should be made in late summer, fall, or winter prior to germination.
(3) Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Winfield Fertilizer with BARRICADE Rate (lb./A)		Months Before Overseeding		
Product	**ai**	**North**	**Transition**	**South**
178	.50	4	4	4
235	.66	5	4	4
264	.74	6	5	5
285	.80	---	6	6
357	1.00	---	7	7
407	1.14	---	---	9
464	1.30	---	---	10
530	1.50	---	---	12

LANDSCAPE ORNAMENTALS
(including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 235 to 530 lbs./A in fall and/or spring. Use higher rates of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 530 lbs./A.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines, and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage), some cultivars of listed plants may be sensitive to this product.

Table 1. Tolerant Ornamental Species-All States

Scientific Name	Common name
*Abies spp.***	Fir species** (Balsam, Fraser, Noble, etc.)
Acer palmatum	Japanese Maple
A. platanoides	Norway Maple
*Actinidia chinensis**	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Aucuba japonica	Japanese Aucuba
Berberis gladwynensis	Barberry
B. julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. thunbergii	Japanese Barberry
B. verruculosa	Warty Barberry
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
*Citrus spp.**	Citrus species*
Cornus florida	Flowering Dogwood
C. stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. microphyllus	Rockspray Cotoneaster
Cretaegus spp.	Hawthorne
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Dodonea viscosa	Hop Bush
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Japanese Spindle Tree (Evergreen Euonymus)
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia spp.	Gaillardia, Blanket Flower: "Goblin"
Gardenia jasminoides	Gardenia, Cape-Jasmine
*Gladiolus spp.***	Gladiolus species**
Hedera helix	English Ivy
*Hibiscus***	Rose of Sharon**
Hibiscus rosa-sinensis	Chinese Hibiscus
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. pernyi	Holly
I. vomitoria	Yaupon Holly
Inula ensifolia	
*Iris spp.***	Iris species**
Jasminium nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper
J. conferta	Shore Juniper: Blue Pacific
J. horizontalis	Creeping Juniper
*Juglans spp.**	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crape Myrtle
Ligustrum amurense	Amur Privet
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Liriope muscari	Big Blue Lilyturf
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
*Magnolia spp.***	Magnolia species**
Maleophora luteola	Ice Plant
*Malus spp.**	Crabapple*
Nandina domestica	Heavenly Bamboo
*Narcissus spp.***	Narcissus species**
Nerium oleander	Oleander
*Olea europaea**	Olive*
Ophiopogon japonicus	Mondo Grass
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
*Persea americana**	Avocado*
Photinia fraseri	Frasier's Photinia (Redtip)
*Picea spp.***	Spruce species** (Colorado Blue, Norway, etc.)
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine
P. canariensis	Canary Island Pine
P. elliottii	Slash Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Black Pine
P. palustrus	Longleaf Pine
P. radiata	Monterey Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
P. taeda	Loblolly Pine
P. virginiana	Virginia Pine
*Pistachio spp.**	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum
P. tobira	Japanese Pittosporum
Podocarpus macrophyllus	Japanese Yew
Prunus laurocerasus	English Laurel
*Prunus spp.**	Almond, Apricot, Nectarine Peach, Plum and Prune*
*Pseudotsuga menziesii***	Douglas Fir**
Pyracantha coccinea	Firethorn (Scarlet)
P. fortuneana	Firethorn
P. koidzumii	Firethorn
Pyrus spp.	Bradford Pear sp.
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorne
Rhododendron (including Azalea)	'Coral Bells', 'Formosa', 'PJM','Roseum Elegans'
Rosa banksine	Lady Bank's Rose
Rosmarinus officinalis'	Rosemary*
Santolina virens	Santolina
Sedum album	Stonecrop
Syzygium paniculatum	Japanese Boxcherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Thuja occidentalis	American Arborvitae
Trachelospermum asiatum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush
V. wrightii	Leatherleaf Viburnum
Vinca major	Vinca
Vinca minor	Dwarf Periwinkle
*Vitis spp.**	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca
Y. filamentosa	Adam's Needle, Yucca

*Do not use on food-producing trees, vines, or plants.
**Not for use on container-grown plants.

Table 2. Tolerant Ornamental Species/Varieties - All States Except CA

Scientific Name	Common name
Abelia grandiflora	Abelia: Sherwood
Achilles spp.	Yarrow: King Edward
Agapanthus orientalis	
Akebia quintara	Five-Leaf or Chocolate Vine
Allium cernuum	Lady's Leek, Nodding Onion
Anemone hybrida	Japanese Anemone
Aquilegia spp.	Aquilegia: Red and Gold
Artemisia spp.	Wormwood: Silver Mound, Castle
Aster spp.	Aster: Bonny Blue, Purple Dome
Aster X frikartii	
Athyrium filiz-femina	Lady Fern: Fem Lady
Begonia spp.	Fibrous Begonia: Hardy Grandis
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Bougainvillea spp.	Bougainvillea
Buddleia davidii	Butterfly-Bush (Dwarf Blue): Royal Red
Caillstemon citrinus	Crimson Bottlebrush
Campanula carpatica	Tussock Bellflower, (White Clips)
Campis X Tagliabuana	Trumpet Creeper, Trumpet Flower, Madame Galen
Ceratostigma plumbagonoides	
Chrysanthemum nipponicum	
Coreopsis spp.	Coreopsis (Calliopsi): Early Sunrise,Moonbeam
Crocosmia spo.	Lucifer
Delosperma spp.	Cooperi Pink
Delphinium spp.	Larkspur: Blue Elf
Dianthus deltoides	Dianthus: Maiden Pinks "Zing"
D. gratuabioik:utabys	Cheddar Pink
Echinacea pupurea	Coneflower, Purple: Magnus
Forsythia suspense	Weeping Forsythia
Gaillardia spp.	Gaillardia, Blanket Flower: "Goblin"
Gaura spp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
Gypsophila repens	Baby's Breath
Hellanthemum spp.	Sunrose
Hemerocallis spp.	Daylily: Aztec Gold, Stella De Oro, Tender Love
Heucherella sop.	Caral Bell: Bridget Bloom
Hibiscus spp.	Mallow: Disco Belle White
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. Sieboldiana	Hosta, "Searsucker"
Houttuytnia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
Inula ensifolia	
Iris ensata	Sword-Leaved Iris: Jodlesong
Iris siberica	Siberian Iris: Cabernet
Juniparus davurica	Parsoni
Lagerstromia indica X Fauriei	Crape Myrtle: Tuscarora
Lantana montevidensis	Weeping Lantana
Lavender spp.	Lavender: Munstead
Leontopodium alpinum	Edelweiss
Ligustrum sinense	Chinese Privet Variegata
Lilium spp.	Lil: Jazz
Liriope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink
Loropetalum chineense	Burgundy
Lythrum spp.	Loosestrife: Modern Pink
Miscanthus sinensis	Yaku Jima, Silberfeder
Oenothera missourensis	Evening Primrose
Osmanthus heterphylius	Osmanthus (False Holly): Gulf Tide
Paeonia suffruticosa	Tree Peony
Pennisetum setaceumi	Fountain Grass (Dwarf)
Perovskia atriplicfolia	
Physotegia veginiana	Dragonhead, False: Vivid
Quercus Shumardii	Oak, Shumard's Red
Raphiolepsis umbellate	Yedda Hawthorn
Rhododendron (including Azalea)	'Delaware Valley White', 'Flame Creeper', 'Girard Crimson', George L. Tabor', 'Wakeiebisu', 'White Gumpo'
Rudbeckia spp.	Black-Eyed Susan: Goldstrum
Saxifraga spp.	Saxifrage: Purple Dome
Scabiosa spp.	Pincushion Flower
Sedum cauticola	Stonecrop: Lidakense
Sedum dasyphyllum	Stonecrop
Sedum spurium	Stonecrop: Dragon's Blood
Spiraea Bumalda	Spirea: Anthony Waterer
Syzygium pariculatum	Australian Brushcherry
Teucrium spp.	Germander
Thalictrum dipterocarpum	Meadow Rue
Veroniac spp.	Veronica, Speedwell: Sunny Border
Viburnum susbensum	Arrowood Viburnum

WARRANTY DISCLAIMER

Winfield Solutions, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

The directions for use of this product must be followed carefully. EXCEPT AS PROVIDED ABOVE AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Distributed by:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589