

ACCEPTED
VIA NOTIFICATION
LABEL NOT REVIEWED

April 25, 2019

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Doc id: 561408

TRIATHLON® BA

Aqueous Suspension Biofungicide/Bactericide

FOR ORGANIC PRODUCTION

Active Ingredient:

Bacillus amyloliquifaciens strain D747* 98.85%

Other Ingredients: 1.15%

Total: 100.00%

*Contains a minimum of 1×10^{10} colony-forming units (cfu) per milliliter

EPA Reg. No. 70051-107-59807

EPA Est. No. 70051-CA-001



For Organic Use

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Inside Panels for Additional Precautionary Statements.

988544

Net Contents: 1 Gallon (3.78 Liters)



8 96531 00249 7

Produced for:
OHP, Inc.
PO Box 746
Bluffton, SC 29910-0746
(800) 356-4647

ESL061118 REV031419



Pantone 288



Pantone 376



Pantone 3295



Black

FIRST AID

IF ON SKIN: Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product label with you when calling a poison control center or doctor. Hot Line No.: 1-800-356-4647 for additional information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco and using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Mixer/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no instructions are available, use detergent and hot water for washables. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides, the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For Terrestrial Uses: Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from treated areas.

GENERAL INFORMATION

TRIATHLON BA is a broad-spectrum preventative biofungicide/bactericide for control or suppression of fungal and bacterial plant diseases. The active ingredient of **TRIATHLON BA** is a naturally occurring strain (D747) of the beneficial bacterium *Bacillus amyloliquefaciens*. **TRIATHLON BA** also colonizes plant root hairs, preventing establishment of disease-causing fungi and bacteria.

TRIATHLON BA can be applied alone or in combination and/or rotation with chemical fungicides as a tool for integrated disease management in agricultural crops, ornamental and nursery plants, and turfgrass. **TRIATHLON BA** offers a valuable tool for management of resistance to chemical fungicides through its multiple and unique modes of action.

TRIATHLON BA can be applied up to and including the day of harvest.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, waterproof gloves, shoes plus socks.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of treated areas until sprays have dried.

MIXING AND HANDLING INSTRUCTIONS

Mix the required amount of **TRIATHLON BA** in water with sufficient agitation to maintain a uniform suspension in the spray or mixing tank. Tank should be cleaned prior to use. Do not use highly alkaline or highly acidic water to mix sprays. Use a buffering agent if necessary to maintain neutrality (pH 6 to 8) of water in the tank. Maintain agitation during application. Apply immediately after mixing; do not allow spray mix to stand overnight.

TRIATHLON BA can be mixed and used with other agricultural chemicals for which such mixing is permitted by the product labels, in accordance with the most restrictive of those label limitations and precautions. If such a mixture is planned, a com-

patibility "jar test" should first be conducted by mixing the correct proportions of **TRIATHLON BA** and these products in a small volume of water.

APPLICATION METHODS

Ground: TRIATHLON BA can be applied in most commonly-used ground application equipment, such as (but not limited to): tractor-mounted boom, airstart, high clearance, hose-end, backpack, and other pressurized sprayers; hose-end or hand-held sprayers; foggers or mist blowers; water wheel and other drench applicators; and shank or other soil injection method.

Aerial: TRIATHLON BA can be applied by fixed or rotary winged aircraft in a minimum of 3 gallons of water per acre. Standard precautions should be taken to minimize spray drift.

Chemigation: TRIATHLON BA can be applied through drip (trickle) and sprinkler type irrigation equipment. Refer to the section entitled "Chemigation Instructions" for detailed instructions.

CROPS	DISEASES/PATHOGENS (See footnotes for additional information)
Vegetables and melons:	
Brassica vegetables such as broccoli, cabbage, cauliflower, Brussels sprouts, kohlrabi, and other cole crops. (including those grown for seed production).	Pin rot complex (<i>Alternaria/Xanthomonas</i>)* Leaf spots (<i>Alternaria</i> spp., <i>Xanthomonas</i> spp.) Downy mildew (<i>Peronospora</i> spp.) Powdery mildew (<i>Erysiphe polygoni</i>) "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , <i>Phytophthora</i> , or <i>Verticillium</i> * spp. (see instructions below for "Soil application").
Bulb vegetables such as onions, garlic, shallots, and others. (including those grown for seed production).	<i>Botrytis</i> spp. (neck rot, leaf blight) Purple blotch (<i>Alternaria</i> spp.) Downy mildew (<i>Peronospora</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) Rust (<i>Puccinia porii</i>)* "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , <i>Phytophthora</i> , or <i>Verticillium</i> * spp. (see instructions below for "Soil application").

CROPS	DISEASES/PATHOGENS (See footnotes for additional information)
Vegetables and melons: (continued)	
Cucurbits such as cucumbers, squash (all types), cantaloupes, muskmelons, watermelons, and other melons. (including those grown for seed production).	<p>Powdery mildew (<i>Erysiphe</i> and <i>Sphaerotheca</i> spp.) Downy mildew (<i>Pseudoperonospora</i> spp.) Gummy stem blight (<i>Didymella bryoniae</i> and <i>Phoma cucurbitacearum</i>)</p> <p>See instructions below for "Soil application" against the following diseases:</p> <p>Vine decline (<i>Monosporascus cannonballus</i>)** Charcoal rot (<i>Macrophomina phaseoli</i>)**</p> <p>"Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i>, <i>Rhizoctonia</i>, <i>Fusarium</i>, <i>Phytophthora</i>, or <i>Verticillium</i>* spp.</p>
Fruiting vegetables such as tomatoes, peppers, eggplant, tomatillo, okra, and others. (including those grown for seed production).	<p>Bacterial spot (<i>Xanthomonas</i> spp.)¹ Bacterial speck (<i>Pseudomonas syringae</i> pv. <i>tomato</i>)¹ Gray mold (<i>Botrytis cinerea</i>) Powdery mildew* (<i>Leveillula</i>, <i>Didiopsis</i>, <i>Erysiphe</i>, and <i>Sphaerotheca</i> spp.) Early blight (<i>Alternaria solani</i>)* Late blight (<i>Phytophthora infestans</i>)*</p> <p>See instructions below for "Soil application" against the following diseases:</p> <p>"Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i>, <i>Rhizoctonia</i>, <i>Fusarium</i>, <i>Phytophthora</i>, or <i>Verticillium</i>* spp.</p> <p>Southern blight (<i>Sclerotium rolfsii</i>)* and **</p>

CROPS	DISEASES/PATHOGENS (See footnotes for additional information)
Vegetables and melons: (continued)	
Leafy vegetables such as head and leaf lettuce, celery, spinach, radicchio, arugula, watercress, and others (including leafy <i>Brassica</i> vegetables such as mustard and collard greens, kale, bok choy, and related crops), including those grown for seed production.	<p>Downy mildew (<i>Bremia lactucae</i>, <i>Peronospora</i> spp.)² Powdery mildew (<i>Golovinomyces</i> (<i>Erysiphe</i>) <i>cichoracearum</i>)* Bacterial blights Head and leaf drop (<i>Sclerotinia</i> spp.)² Pink rot (<i>Sclerotinia sclerotiorum</i>)² Leaf spots (<i>Cercospora</i> spp.)</p> <p>See instructions below for "Soil application" against the following diseases:</p> <p>"Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i>, <i>Rhizoctonia</i>, <i>Fusarium</i>, <i>Phytophthora</i>, or <i>Verticillium</i>* spp.</p> <p>Bottom rot (<i>Rhizoctonia solani</i>)</p>
Legume vegetables succulent and dried beans and peas such as green, snap, shell, and Lima beans, garbanzo beans, chickpeas, soybeans, dry beans, peas, split peas, lentils, and other legumes, including those grown for seed production.	<p>White mold (<i>Sclerotinia sclerotiorum</i>)² Gray mold (<i>Botrytis cinerea</i>) Powdery mildew (<i>Microsphaera diffusa</i>) Rusts*, including <i>Uromyces appendiculatus</i>, <i>Puccinia</i> spp., and Asian soybean rust (<i>Phyospora pachyrhizi</i>)</p> <p>"Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i>, <i>Rhizoctonia</i>, <i>Fusarium</i>, <i>Phytophthora</i>, or <i>Verticillium</i>* spp. (see instructions below for "Soil application").</p>

CROPS	DISEASES/PATHOGENS (See footnotes for additional information)
Vegetables and melons: <i>(continued)</i>	
Root, tuber, and corn vegetables such as potato, sweet potato, carrot, cassava, beets, ginger, radish, horseradish ²² , ginseng, turnip, and other root, tuber and corn crops. (including those grown for seed production).	Black root/crown rot (<i>Alternaria</i> spp.) Bacterial leaf blight (<i>Xanthomonas campestris</i>) Downy mildew (<i>Peronospora</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) Gray mold (<i>Botrytis</i> spp.) White mold (<i>Sclerotinia sclerotiorum</i>) [†] Black leg/bacterial soft rot (<i>Erwinia carotovora</i>)** Early blight (<i>Alternaria solani</i>) [*] Late blight (<i>Phytophthora infestans</i>) [*] See instructions below for "Soil application" against the following diseases: Black scurf (<i>Rhizoctonia solani</i>) Cavity spot (<i>Pythium</i> spp.) "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , <i>Phytophthora</i> , or <i>Verticillium</i> * spp.
Other vegetables such as sweet corn, popcorn, asparagus, peanut, and watercress	<i>Botrytis</i> spp. Rusts (<i>Puccinia</i> spp.) White mold (<i>Sclerotinia sclerotiorum</i>) [†] Leaf spots (<i>Cercospora</i> and <i>Cercosporidium</i> spp.) [*] "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , <i>Phytophthora</i> , or <i>Verticillium</i> * spp. (see instructions below for "Soil application").

CROPS	DISEASES/PATHOGENS (See footnotes for additional information)
Tree fruits and nuts	
Citrus such as orange, lemon, lime, grapefruit, tangerine (mandarin), tangelo, pummelo, and other citrus	<i>Alternaria</i> leaf spot (<i>Alternaria alternata</i>) Postbloom fruit drop (<i>Colletotrichum acutatum</i>) [*] Greasy spot (<i>Mycosphaerella citri</i>) ^{6,8} Citrus canker (<i>Xanthomonas campestris</i> pv. <i>citri</i>) [*] Scab (<i>Elsinoe fawcettii</i>) ^{4,8} Melanose (<i>Diaporthe citri</i>) [*]
Pome fruits such as apple, pear, crabapple, quince, and others	Powdery mildew (<i>Podosphaera leucotricha</i>) [†] Scab (<i>Venturia</i> spp.) [*] Flyspeck (<i>Zygophiala jamaicensis</i>)** Sooty blotch disease complex ^{6,8,9} Brooks spot (<i>Mycosphaerella pomii</i>)*** Bot rot/white rot (<i>Botryosphaeria dothidea</i>) ^{8,9,10} Bitter rot (<i>Colletotrichum</i> spp.) ⁹ Cedar apple rust (<i>Gymnosporangium juniperi-virginianae</i>)*** Fire blight (<i>Erwinia amylovora</i>) ⁷
Stone fruits such as apricot, cherry, nectarine, peach, plum, prune, pluot, and others	Powdery mildew (<i>Sphaerotheca</i> and <i>Podosphaera</i> spp.) ^{8,9} Bacterial canker (<i>Pseudomonas</i> spp.) Brown rot blossom blight (<i>Monilinia laxa</i>) ⁹ Brown rot (<i>Monilinia fructicola</i>) ^{8,10} Gray mold (<i>Botrytis cinerea</i>) ⁹ Peach leaf curl (<i>Taphrina deformans</i>) Bacterial leaf spot (<i>Xanthomonas arboricola</i> pv. <i>prunij</i>) [*] Rusty spot (<i>Podosphaera leucotricha</i>) [†]
Tree nuts such as almond, pistachio, pecan, walnut, filbert, hazelnut, chestnut, macadamia, and other tree nuts.	Walnut blight (<i>Xanthomonas campestris</i>) ¹¹ Anthracnose (<i>Colletotrichum acutatum</i>) [*] Bacterial canker (<i>Pseudomonas syringae</i>) [*] Shot hole (<i>Wilsonomyces carpophilus</i>) [*] Brown rot (<i>Monilinia</i> spp.) [*] Pecan scab (<i>Cladosporium caryigenum</i>) ¹ and **

CROPS	DISEASES/PATHOGENS (See footnotes for additional information)
Tree fruits and nuts <i>(continued)</i>	
Pomegranates	Leaf and fruit spots (<i>Cercospora</i> , <i>Gloeosporium</i> and <i>Pestalotia</i> spp.) ¹ Fruit rots (<i>Alternaria</i> , <i>Botrytis</i> , and other spp.) ¹⁰ Powdery mildew (<i>Sphaerotheca pannosa</i>)
Other fruits	
Strawberry	Powdery mildew (<i>Sphaerotheca macularis</i> , <i>Erysiphe</i> spp.) ^{*12} Gray mold (<i>Botrytis cinerea</i>) ^{*11} Anthracnose (<i>Colletotrichum acutatum</i>) Angular leaf spot (<i>Xanthomonas fragariae</i>) ¹ For the following diseases, see instructions below for "Soil application" (and also root dip instructions ²⁰): "Damping off" and root or crown diseases caused by <i>Rhizoctonia</i> , <i>Fusarium</i> , <i>Pythium</i> , <i>Phytophthora</i> , and/or <i>Verticillium</i> * spp. Charcoal rot (<i>Macrophomina phaseolina</i>) ^{**}
Berries , including blueberry, blackberry, raspberry, loganberry, huckleberry, kiwifruit, gooseberry, elderberry, cranberry (non-flooded fields), currant, and other berries	Mummy berry (<i>Monilinia vaccinii-corymbosi</i>) [*] Botrytis blight (<i>Botrytis cinerea</i>) Bacterial canker (<i>Pseudomonas</i> spp.) ¹³ Anthracnose fruit rot (<i>Colletotrichum acutatum</i>) ¹⁰ Sclerotinia (<i>Sclerotinia sclerotiorum</i>)
Grapes including wine grapes, table grapes, and raisins	Powdery mildew (<i>Erysiphe</i> (formerly <i>Uncinula</i>) <i>necator</i>) ¹⁴ Gray mold (<i>Botrytis cinerea</i>) ¹⁵ Sour rot complex ¹⁵ Downy mildew (<i>Plasmopara viticola</i>) [*] Phomopsis (<i>Phomopsis viticola</i>) ¹⁶ Eutypa (<i>Eutypa lata</i>) ¹⁷

CROPS	DISEASES/PATHOGENS (See footnotes for additional information)
Other fruits <i>(continued)</i>	
Tropical fruits such as avocado ¹⁸ , mango ¹⁸ , papaya ¹⁸ , pineapple ¹⁹ , banana, plantain, and others.	Anthracnose (<i>Colletotrichum</i> spp.) Scab (<i>Sphaceloma perseeae</i>) Bacterial canker (<i>Xanthomonas campestris</i>) Sigatoka (<i>Mycosphaerella fijiensis</i>) ²⁰
Other Crops	
Herbs and spices such as basil, thyme, coriander, dill, cilantro, parsley, mint, and others.	Powdery mildews (<i>Oidium</i> spp. and others) Downy mildews (<i>Peronospora</i> spp. and others) [*] Damping off diseases (<i>Rhizoctonia</i> , <i>Pythium</i> , <i>Alternaria</i> , and <i>Fusarium</i> spp.) Leaf spots (<i>Alternaria</i> , <i>Septoria</i> , <i>Colletotrichum</i> , and <i>Cercospora</i> spp.) [*] Bacterial diseases (<i>Erwinia</i> , <i>Xanthomonas</i> , and <i>Pseudomonas</i> spp.) Rusts (<i>Puccinia</i> spp. and others) "Damping off" and root or crown diseases caused by <i>Rhizoctonia</i> , <i>Fusarium</i> , <i>Pythium</i> , <i>Phytophthora</i> , and/or <i>Verticillium</i> * spp. (see instructions below for "Soil application").
Footnotes:	
* Suppression only; for improved control mix or rotate with chemical fungicide approved for such use. **NOT FOR USE IN CALIFORNIA	
¹ Tank mix or rotate with copper-based fungicides at label rates for improved control.	
² Apply at or immediately following planting (but before plant emergence) as a banded seedline treatment 4 to 6 inches wide. Make second application at thinning or cultivation in sufficient water and multiple nozzles to ensure thorough coverage of lower leaves and surrounding soil surface. Incorporation with light irrigation after application may improve disease control. Repeat at 10-14 day intervals if conditions promoting disease persist.	
³ For greasy spot suppression, apply at first new foliar flush and repeat with each new flush. Tank mix with spray oil or copper based fungicide at labeled rates.	

Footnotes: (continued)

- ⁴For suppression of citrus scab, start applications at first new foliage flush and repeat at petal fall and when fruit are 1/2 inch in diameter.
- ⁵Make first application at or before tight cluster if conditions favor disease development. Repeat at 7-10 day intervals through the second cover spray or longer on susceptible varieties or if environmental conditions favor rapid disease development.
- ⁶Begin applications before bloom when environmental conditions favor disease development, repeating at 7 to 14 day intervals or as needed. Control may be enhanced by addition of a surfactant to improve spray coverage. Use only surfactants known to be safe for use on the crop and for which such use is allowed.
- ⁷Rotate with antibiotics registered for fire blight control for improved performance. Begin applications at 1-5% open blossoms and repeat every 3-7 days as necessary until petal fall, when intervals can be increased to 7 days. **TRIATHLON BA** can also be used in summer "cover spray" applications to control the shoot blight phase of fire blight and summer diseases. Can be mixed with copper fungicides to improve control.
- ⁸Make first application at popcorn stage and repeat every 7 days.
- ⁹Start applying at early bloom stage and repeat every 7 days through petal fall.
- ¹⁰Pre-harvest applications in sufficient water to cover fruit or other harvested plant parts may improve control of postharvest infections.
- ¹¹Begin applications at or before pistillate bloom, repeating every 7-10 days. Apply before rainfall if possible, and tank mix or rotate with a copper-based bactericide registered for such use for improved control.
- ¹²Start applications at or just before flowering and repeat every 7-10 days as needed through harvest.
- ¹³Apply before fall rains and again during dormancy before spring growth.
- ¹⁴Start applications when new shoots are 1/2 to 1 1/2 inches long. Repeat at 3-5 inches, 8-10 inches, and then at 7-10 day intervals until disease conditions no longer exist.
- ¹⁵Apply at bloom, before bunch closure, at veraison, and before harvest.
- ¹⁶Apply when shoots are 1/2 to 1 inch long and again when 6-8 inches long.
- ¹⁷Mix 2 fluid ounces **TRIATHLON BA** per gallon of water and apply to pruning wounds.
- ¹⁸Apply at budbreak and repeat on 14-21 day interval as needed through harvest.
- ¹⁹Apply at flowering and repeat on 14-21 day interval as needed through harvest.

Footnotes: (continued)

- ²⁰Apply at first appearance of leaves and repeat at 7-21 day intervals as needed, in sufficient water to obtain thorough coverage of foliage. Tank mix with spray oil or other registered fungicides for improved control.
- ²¹Mix 6 to 10 fluid ounces **TRIATHLON BA** per 100 gallons of water and apply in minimum of 20 gallons per acre from emergence to training, 50 gallons per acre from training to wire, and 100 gallons per acre from wire touch through harvest.
- ²²For treatment of horseradish or strawberry roots immediately before transplanting: immerse bare roots (individually or in bunches) for 10 seconds in a suspension of 1 to 2 pints **TRIATHLON BA** per gallon of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PEST or DISEASE:

For suppression of root and collar rots caused by *Phytophthora*, *Pythium*, *Fusarium*, *Rhizoctonia*, *Armillaria*, and other soilborne pathogens in bearing and non-bearing tree, nut, and vine crops, including:

Citrus:	Orange, Grapefruit, Lemon, Lime, Pummelo, Tangerine (Mandarin), Tangelo, and other citrus;
Grapevines:	Wine grapes, Table grapes, and Raisins;
Pome fruit:	Apple, Pear, Crabapple, Quince, and other pome fruits;
Stone fruit:	Apricot, Cherry, Nectarine, Peach, Plum, Prune, <i>Prunus</i> hybrids (such as Pluot, Aprium, Apriplum, and Plumcot), and other stone fruit;
Tree nuts:	Almond, Chestnut, Pistachio, Pecan, Walnut, Hazelnut (Filbert), Macadamia and other tree nuts;
Other tree fruits:	Avocado, Mango, Papaya, and Pomegranate.

APPLICATION INSTRUCTIONS:

For root diseases: Apply 1 to 2 quarts of **TRIATHLON BA** per acre as a banded soil spray or drench, or as chemigated injection via microsprinkler, drip (elevated, buried or ground-lay) or other irrigation systems. Apply in sufficient water or irrigate immediately after application to move the product to the root zone. Begin applications in early spring, timed for root flush and early shoot growth. Continue applications at 4-6 week intervals through fall root flush.



For collar rots: Apply as drench or spray at the base of the trunk, covering the soil contact zone.

Foliar application: For control of diseases on foliage, flowers, fruit, or other above-ground parts of plants: Mix **TRIATHLON BA** in water and apply as a spray at a rate of **0.5 to 6 quarts of TRIATHLON BA** per acre in sufficient water to achieve thorough coverage of the crop canopy with minimal runoff. Begin applications at crop emergence, transplanting, or when conditions are conducive to development of disease. Repeat application every 3 to 10 days, or as needed, for as long as conditions favor disease development. Lower rates (0.5 to 3 quarts per acre) may be applied under light disease pressure, to smaller (e.g. newly-emerged) plants, or when **TRIATHLON BA** is used in a tank mix with other fungicides whose labels allow such use. Under moderate to severe disease pressure, or when environmental conditions and plant stage are conducive to rapid disease development, use higher label rates (3-6 quarts/acre), apply more frequently (every 3-7 days), and mix or rotate **TRIATHLON BA** with other fungicides for improved performance.

Soil application: For control of soilborne diseases infecting seeds, seedlings, roots, crown, stems, or other plant parts below ground or in contact with soil: Apply **TRIATHLON BA** at **0.5 to 4.5 pints per acre**. Mix the required amount in sufficient water to apply by one of the following methods:

- Soil drench applied to transplants in flats or pots in the greenhouse or nursery any time prior to transplanting (see additional drench instructions under "Nurseries, greenhouses, shade houses, and ornamental plants" below).
- Soil drench at transplanting, using a "water wheel" injector, spray nozzles/hoses, or other method to drench each root ball and/or planting hole.

Follow-up (post-planting) preventative applications can be made every 2-4 weeks by one or more of the following methods, if needed:

- Drip (trickle) or any type of sprinkler irrigation, any time after planting or transplanting. See Chemigation Instructions for additional information.
- Spray directly onto the soil surface and/or lower plant parts. If targeting root disease, follow immediately with sufficient overhead sprinkler irrigation to move **TRIATHLON BA** to the root zone.
- Injection directly into the rooting zone using shanks or similar equipment.

Lower rates (0.5 to 2 pints of **TRIATHLON BA** per acre) may be applied under light disease pressure, to smaller plants, or when **TRIATHLON BA** is used in a tank mix with other fungicides whose labels allow such use. Under moderate to severe disease pressure, or when environmental conditions and plant stage are conducive to rapid disease development, use higher label rates (2 - 4.5 pints per acre), apply

more frequently (every 2 weeks), and mix or rotate **TRIATHLON BA** with other fungicides for improved performance.

Nurseries, greenhouses, shadehouses, and ornamental plants

Spray application: Mix **0.5 to 6 quarts of TRIATHLON BA** per 100 gallons of **water** and apply as a foliar spray of sufficient volume to wet the entire plant with minimal runoff. Begin preventative applications at plant emergence and repeat every 3-28 days as needed (every 3-7 days if disease pressure is high or environmental conditions are highly favorable to disease outbreak, 10-28 days under low pressure or less conducive conditions).

Drench application: Mix **0.5 to 4.5 pints of TRIATHLON BA** per 100 gallons of **water** and apply as a drench or coarse spray to soil or other growing media in pots, flats, plugs, trays, or planting beds, for control or suppression of soilborne diseases of seedlings, cuttings, bedding plants, and transplants (including vegetables and other transplanted food crops). Make first application at or immediately before seeding, sticking, germination, or transplanting. Repeat applications every 14-28 days as needed. Transplants can be treated immediately before transplanting into field soils to protect against damping-off and other diseases that reduce plant establishment.

Cutting or root dip: Dip basal end of cuttings or bare roots (individually or in bunches) in a suspension of **1 to 2 pints of TRIATHLON BA** per gallon of **water**. Immerse for 5-10 seconds immediately before planting.

Chemigation: Mix **0.5 to 4.5 pints of TRIATHLON BA** per 100 gallons of **water** and apply via drip, handheld, or sprinkler irrigation systems. Refer to "Chemigation Instructions" for more details.



CROPS/USE SITES	DISEASES/PATHOGENS
Indoor, outdoor, and shade- or other cover-grown ornamental trees and shrubs, flowering plants, foliage plants, tropical plants, potted plants, potted or cut flowers, bedding plants, forestry seedlings, conifer production for reforestation, fruit trees, vegetables and other crops grown in greenhouses or nurseries, or other cover, interiorscapes, and landscapes.	<p>Powdery mildews caused by <i>Erysiphe</i>, <i>Podosphaera</i>, <i>Sphaerotheca</i>, <i>Oidium</i>, and <i>Golovinomyces</i> spp.</p> <p>Anthracnose (<i>Colletotrichum</i> spp.)</p> <p>Bacterial leaf spots caused by <i>Erwinia</i>, <i>Pseudomonas</i>, and <i>Xanthomonas</i> spp.</p> <p>Damping-off disease (<i>Rhizoctonia</i>, <i>Pythium</i>, <i>Fusarium</i> spp.)</p> <p>Late blight, blackeye, and root rots caused by <i>Phytophthora</i> spp.</p> <p>Gray mold and blight caused by <i>Botrytis cinerea</i></p> <p>Black root rot (<i>Aspergillus</i> spp.)</p> <p>Black spot of roses (<i>Diplocarpon rosae</i>)</p> <p>Downy mildew (<i>Peronospora</i> spp.)</p> <p>Leaf spots caused by <i>Alternaria</i>, <i>Septoria</i>, <i>Cercospora</i>, <i>Entomosporium</i>, <i>Helminthosporium</i>, and <i>Myrothecium</i> spp.</p> <p>Rust (<i>Puccinia</i> spp.)</p> <p>Scab (<i>Venturia</i> spp.)</p> <p>Root rot, bottom rot, or stem rot caused by <i>Rhizoctonia solani</i></p> <p><i>Sclerotinia</i> blight</p> <p><i>Fusarium</i> wilts</p>

Turfgrass application

For control of foliar diseases, apply **TRIATHLON BA** at **1 to 4 fluid ounces per 1,000 square feet** as a ground-directed spray in sufficient water to provide thorough coverage. To control root and crown diseases in or on the soil, immediately follow the spray with sufficient overhead sprinkler irrigation to move the product into the root zone.

USE SITES/CROPS	DISEASES/PATHOGENS
Turf, sod, lawns, golf course (fairways, roughs, greens, tees), grass seed production	<p>Anthracnose (<i>Colletotrichum graminicola</i>)</p> <p>Brown patch (<i>Rhizoctonia solani</i>)</p> <p>Dollar spot</p> <p>Including but not limited to: Bluegrass, Bentgrass, Bermudagrass (common & hybrid), Dichondra, Fescue, Orchardgrass, <i>Poa annua</i>, St. Augustine grass, Ryegrass, <i>Zoysia</i>, mixtures, and other grasses or ornamental turf</p>
	<p><i>Lanzia</i> and <i>Moellerodiscus</i> spp., formerly <i>Sclerotinia homeocarpa</i>)</p> <p>Powdery mildew (<i>Erysiphe graminis</i>)</p> <p>Rust (<i>Puccinia</i> spp.)</p> <p>Gray leaf spot (<i>Pyricularia grisea</i>)</p> <p>"Damping off" or seedling blights caused by <i>Pythium</i></p>

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a dry area inaccessible to children. Store in original containers only. Keep container closed when not in use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. **-for containers equal to or less than 5 gallons-**

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CHEMIGATION INSTRUCTIONS

General information:

Apply this product only through drip (trickle) irrigation (including micro-irrigation through spaghetti tubes or individual tubes) or sprinkler irrigation (including impact or microsprinklers, microjet, overhead boom, water gun, solid set, lateral move, end tow, side-roll, center pivot, or hand move, including mist-type systems); or with hand-held calibrated irrigation equipment (such as a hand-held wand with injector). Do not apply this product through any other type of irrigation system.

Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where

there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment. Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and injector system and flush with clean water before use. Failure to provide a clean tank, free of scale or residues may reduce effectiveness of this product.

Drip (trickle) and micro-irrigation chemigation

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (i.e., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Dilute the product in water following the label mixing directions. It may be pre-mixed in a supply tank with water, fertilizer, or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff or excessive drip from pots. Application should be continuous in sufficient water to apply the recommended rate evenly to the entire treated area.

Sprinkler chemigation:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (i.e., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Dilute the product in water following the label mixing directions. It may be pre-mixed in a supply tank with water, fertilizer or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff or excessive drip from pots. Application should be continuous in sufficient water to apply the recommended rate evenly to the entire treated area.
8. Do not apply when wind speed favors drift beyond the area intended for treatment.

WARRANTY

OHP, Inc. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purpose referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the crop, incompatibility with other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.



Triathlon is a registered trademark of OHP, Inc.

Produced for:

OHP, Inc.

PO Box 746

Bluffton, SC 29910-0746

(800) 356-4647

ESL061118 REV031419



TRIATHLON® BA

Aqueous Suspension Biofungicide/Bactericide

FOR ORGANIC PRODUCTION

Active Ingredient:

Bacillus amyloliquefaciens strain D747* 98.85%

Other Ingredients: 1.15%

Total: 100.00%

*Contains a minimum of 1×10^{10}
colony-forming units (cfu) per milliliter

EPA Reg. No. 70051-107-59807

EPA Est. No. 70051-CA-001



KEEP OUT OF REACH OF CHILDREN CAUTION

See Inside Panels for Additional Precautionary Statements.

Produced for:

OHP, Inc.

PO Box 746

Bluffton, SC 29910-0746

(800) 356-4647

988544

Net Contents:

1 Gallon (3.78 Liters)



ESL061118 REV031419

FIRST AID - IF ON SKIN: Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product label with you when calling a poison control center or doctor. Hot Line No.: 1-800-356-4647 for additional information.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS & DOMESTIC ANIMALS - CAUTION. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco and using the toilet.

ENVIRONMENTAL HAZARDS - For Terrestrial Uses: Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from treated areas.

DIRECTIONS FOR USE - It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **Pesticide Storage:** Store in a dry area inaccessible to children. Store in original containers only. Keep container closed when not in use. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility. **Container Handling:** Nonrefillable container. Do not reuse or refill this container. **-for containers equal to or less than 5 gallons-** Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PULL HERE TO OPEN
PRESS TO RESEAL



Pantone 288



Pantone 376



Pantone 3295



Black

988544 ESL061118 REV031419

